

Separation of variables on a non-hyperelliptic curve

V.G. Marikhin and V.V. Sokolov

February 8, 2008

1 Introduction

In the paper we consider several models that admit a separation of variables on the following algebraic curve of genus 4:

$$\Phi(\xi, Y) = s_6 Y^6 + l(\xi) Y^4 + k(\xi) Y^2 - S(\xi) = 0, \quad (1.1)$$

where

$$s_6 = \frac{\delta^6}{6!} S^{VI}(\xi), \quad k(\xi) = \frac{\delta^2}{10} S'''(\xi) + 4(\alpha \xi^2 + e_2 \xi + e_1), \quad l(\xi) = \frac{\delta^4}{4!} S^{IV}(\xi) - \frac{\delta^2}{2} k''(\xi). \quad (1.2)$$

Here α, δ are parameters, S is arbitrary sixth degree polynomial. If $\delta = 0$, then the curve is hyperelliptic of genus 3.

The class of curves (1.1) with $\delta \neq 0$ can be described as follows. Let $P(\eta, \xi) = 0$ be an arbitrary cubic. In the generic case it is an elliptic curve. Then (1.1) is a double cover over the cubic defined by the formula $\eta = \xi^2 - \delta^2 Y^2$.

To present our approach, we start with the Hamiltonian

$$H = ap_1^2 + cp_2^2 + dp_1 + ep_2 + f, \quad (1.3)$$

where

$$\begin{aligned} a &= -\frac{4s_2 S(s_1)}{s_1 - s_2}, \quad c = \frac{4s_1 S(s_2)}{s_1 - s_2}, \quad d = -s_1 \frac{J}{s_1 - s_2}, \quad e = -s_2 \frac{J}{s_1 - s_2}, \\ f &= \frac{\delta^2}{40} \frac{s_2 S'''(s_1) - s_1 S'''(s_2)}{s_1 - s_2} - \frac{\delta^2}{4} \frac{s_2 S'(s_1) + s_1 S'(s_2)}{(s_1 - s_2)^2} + \\ &\quad \frac{3\delta^2}{4} \frac{s_2 S(s_1) - s_1 S(s_2)}{(s_1 - s_2)^3} + \alpha s_1 s_2, \end{aligned} \quad (1.4)$$

and

$$J = 2\delta \frac{\sqrt{S(s_1)}\sqrt{S(s_2)}}{s_1 - s_2}$$

as the most general model. It can be easily verified that H commutes with the function

$$K = Ap_1^2 + Cp_2^2 + Dp_1 + Ep_2 + F, \quad (1.5)$$

where

$$\begin{aligned} A &= \frac{4S(s_1)}{s_1 - s_2}, & C &= -\frac{4S(s_2)}{s_1 - s_2}, & D &= \frac{J}{s_1 - s_2}, & E &= \frac{J}{s_1 - s_2}, \\ F &= \frac{\delta^2}{40} \frac{S''(s_2) - S''(s_1)}{s_1 - s_2} + \frac{\delta^2}{4} \frac{S'(s_1) + S'(s_2)}{(s_1 - s_2)^2} + \frac{3\delta^2}{4} \frac{S(s_2) - S(s_1)}{(s_1 - s_2)^3} - \alpha(s_1 + s_2) \end{aligned} \quad (1.6)$$

with respect to the standard Poisson bracket $\{p_\alpha, s_\beta\} = \delta_{\alpha\beta}$. If $\delta = 0$, then the terms in H and K linear in momenta disappear and H belongs to the Stäckel class. In this case s_1, s_2 are separated variables. The corresponding Abel's transformation on the hyperelliptic curve of genus 3 has the form

$$\frac{ds_1}{\sqrt{T(s_1)}} + \frac{ds_2}{\sqrt{T(s_2)}} = dt, \quad \frac{s_1 ds_1}{\sqrt{T(s_1)}} + \frac{s_2 ds_2}{\sqrt{T(s_2)}} = 0.$$

Here

$$T(s) = 16 S(s) (\alpha s^2 + e_2 s + e_1),$$

and the constants e_1 and e_2 are values of the integrals H and K , respectively.

The main result of the paper is an explicit formula for the action in the case $\delta \neq 0$ depending on two parameters. We find it directly from the Hamilton-Jacobi equation.

In Sections 4-6 we find the action and a separation of variables for the Kowalewski hyrostat, the Clebsch and the $so(4)$ Schottky-Manakov spinning tops in a way analogous to that used in Sections 2-3 for the Hamiltonian (1.3). Many significant papers (see, for instance, [1, 4, 6, 2, 3, 18, 19, 7, 11, 12, 13, 5, 21] and references there-in) are devoted to these models. In particular, it is known [20, 21]) that the models are linked via some changes of variables. However according to a common opinion of experts, a "satisfactory" separation of variables for the above models is still not found.

For the Clebsch and the $so(4)$ Schottky-Manakov tops S is a polynomial of degree 4. An appearance of the corresponding algebraic curve of genus 3 could be predicted: the characteristic curve for the 3×3 -Lax operator found by A. Perelomov [5] for the Clebsch top has the same form. However in this paper we don't use any Lax representations and the curve arises during computations in a rather natural manner.

The case $S = \text{const}$ also is of interest. The inhomogeneous hydrodynamic type system (2.7) corresponding to the pair (1.3), (1.5) coincides with the Gibbons-Tsarev equation [9].

The separation of variables found in this paper gives rise to a family of elliptic solutions for the Gibbons-Tsarev equation (see Section 7).

Acknowledgments. The authors are grateful to A.V. Borisov, Yu. N. Fedorov, E.V. Ferapontov, V.B. Kuznetsov, and F. Magri for useful discussions. The second author (V.S.) is grateful to the Max Planck Institute for hospitality and financial support. The research was also partially supported by RFBR grants 02-01-00431, and NSh 1716.2003.1.

2 Equations of motion

The Hamilton equations corresponding to (1.3) and (1.5) have the form

$$\frac{ds_1}{dt} = 2ap_1 + d, \quad \frac{ds_2}{dt} = 2cp_2 + e. \quad (2.1)$$

and

$$\frac{ds_1}{d\tau} = 2Ap_1 + D, \quad \frac{ds_2}{d\tau} = 2Cp_2 + E. \quad (2.2)$$

Finding p_1, p_2 from (2.1) and substituting them in the relations $H = e_1$ and $K = e_2$, we get

$$(s_1 - s_2) \left[s_1 \frac{\dot{s}_1^2}{S(s_1)} - s_2 \frac{\dot{s}_2^2}{S(s_2)} \right] + 16(e_1 - f)s_1s_2 + \frac{4\delta^2}{(s_1 - s_2)^3} (s_2^3 S(s_1) - s_1^3 S(s_2)) = 0 \quad (2.3)$$

and

$$(s_1 - s_2)s_1s_2 \left(\frac{\dot{s}_1^2}{S(s_1)} - \frac{\dot{s}_2^2}{S(s_2)} \right) - 4\delta\sqrt{S(s_1)S(s_2)} \left(s_1^2 \frac{\dot{s}_1}{S(s_1)} + s_2^2 \frac{\dot{s}_2}{S(s_2)} \right) + \quad (2.4)$$

$$16s_1s_2((e_1 - f)(s_1 + s_2) + (e_2 - F)s_1s_2) - \frac{4\delta^2}{(s_1 - s_2)^3} (s_2^3 S(s_1)(s_1 - 2s_2) - s_1^3 S(s_2)(s_2 - 2s_1)) = 0,$$

where f and F are defined by (1.4), (1.6).

One of the basic technical problems is a rewriting of this system in a compact form. The straightforward elimination say \dot{s}_2 leads to a cumbersome equation of degree 4 with respect to \dot{s}_1 .

Proposition 1. *Let (u, v) be any solution of the following system of equations:*

$$L(s_2)u^2 - 2L(s_1)v - M(s_2, s_1) = 0, \quad L(s_1)v^2 - 2L(s_2)u - M(s_1, s_2) = 0, \quad (2.5)$$

where

$$M(x, y) = 3L(y) + L'(y)(x - y) + k(y)(x - y)^2, \quad L(x) = \delta^2 S(x).$$

Then the derivatives

$$\dot{s}_1 = -J \frac{s_2 u + s_1}{s_1 - s_2}, \quad \dot{s}_2 = -J \frac{s_1 v + s_2}{s_1 - s_2} \quad (2.6)$$

satisfy the system (2.3), (2.4).

The elimination of variables p_1, p_2 from (2.1), (2.2) leads to the inhomogeneous hydrodynamic type system

$$(s_1)_t + s_2(s_1)_\tau = -J, \quad (s_2)_t + s_1(s_2)_\tau = J. \quad (2.7)$$

It follows from this formula and from (2.6) that

$$(s_1)_\tau = J \frac{u+1}{s_1-s_2}, \quad (s_2)_\tau = J \frac{v+1}{s_1-s_2}. \quad (2.8)$$

Moreover, we find from (2.6) that

$$p_1 = \frac{J u}{8S(s_1)}, \quad p_2 = -\frac{J v}{8S(s_2)}. \quad (2.9)$$

The following parameterization for solutions of (2.5) turns out to be crucial for the separation of variables.

Proposition 2. *Let z_1, z_2, z_3 be solutions of the equation*

$$z^6 L(s_2)^2 - L(s_2)M(s_2, s_1)z^4 + M(s_1, s_2)L(s_1)z^2 - L(s_1)^2 = 0 \quad (2.10)$$

such that

$$\frac{L(x)}{L(y)} = z_1 z_2 z_3, \quad \frac{M(y, x)}{L(y)} = z_1^2 + z_2^2 + z_3^2, \quad \frac{M(x, y)}{L(x)} = \frac{1}{z_1^2} + \frac{1}{z_2^2} + \frac{1}{z_3^2}.$$

Then

$$\begin{aligned} (u_1, v_1) &= (z_1 + z_2 + z_3, \frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3}), & (u_2, v_2) &= (z_1 - z_2 - z_3, \frac{1}{z_1} - \frac{1}{z_2} - \frac{1}{z_3}), \\ (u_3, v_3) &= (z_2 - z_1 - z_3, \frac{1}{z_2} - \frac{1}{z_1} - \frac{1}{z_3}), & (u_4, v_4) &= (z_3 - z_1 - z_2, \frac{1}{z_3} - \frac{1}{z_1} - \frac{1}{z_2}) \end{aligned} \quad (2.11)$$

are solutions of (2.5).

3 Separation of variables.

To separate the variables in the case $\delta \neq 0$, we find the action function $\tilde{S}(s_1, s_2)$ in an explicit form.

Let us consider the system of equations

$$\Phi(\xi, Y) = 0, \quad Y^2 = \frac{1}{\delta^2}(\xi - s_1)(\xi - s_2),$$

where the polynomial $\Phi(\xi, Y)$ is given by (1.1). It is easy to verify that if we substitute to the first equation the expression for Y^2 taken from the second equation, then leading powers of ξ cancel and the equation for ξ turns out to be cubic. By $\xi_i(s_1, s_2)$, $i = 1, 2, 3$ denote the roots of this equation.

Theorem. *The function*

$$\tilde{S}(s_1, s_2, e_1, e_2) = \frac{1}{4} \sum_{n=1}^3 \left[\delta \operatorname{arctanh} \frac{\xi_n - \frac{1}{2}(s_1 + s_2)}{\delta Y(\xi_n)} - \int^{\xi_n} \frac{d\xi}{Y(\xi)} \right] \quad (3.1)$$

satisfies the Hamilton-Jacobi equation

$$H \left(\frac{\partial \tilde{S}}{\partial s_1}, \frac{\partial \tilde{S}}{\partial s_2}, s_1, s_2 \right) = e_1, \quad K \left(\frac{\partial \tilde{S}}{\partial s_1}, \frac{\partial \tilde{S}}{\partial s_2}, s_1, s_2 \right) = e_2,$$

where the functions $H(p_1, p_2, s_1, s_2)$ and $K(p_1, p_2, s_1, s_2)$ are defined by formulas (1.3)-(1.5).

Proof. It follows from (2.9) that the partial derivatives of the action \tilde{S} are given by:

$$\frac{\partial \tilde{S}}{\partial s_1} = \frac{Ju}{8S(s_1)}, \quad \frac{\partial \tilde{S}}{\partial s_2} = -\frac{Jv}{8S(s_2)}. \quad (3.2)$$

The compatibility condition for this system has the form

$$S(s_2) \frac{\partial}{\partial s_2} (Ju) + S(s_1) \frac{\partial}{\partial s_1} (Jv) = 0.$$

The latter relation follows from the identities

$$\begin{aligned} (L'(y) + M_x(x, y))(x - y) + 6L(y) - 2M(x, y) &= 0, \\ (x - y)(M_y(x, y) + M_x(y, x)) + 4(M(x, y) - M(y, x)) &= 0, \end{aligned} \quad (3.3)$$

which can be verified straightforwardly.

Consider an auxiliary Hamiltonian of the form (compare with formula (2.10))

$$\tilde{H} = p_1^3 L(s_1) + p_1^2 p_2 M(s_2, s_1) + p_1 p_2^2 M(s_1, s_2) + p_2^3 L(s_2). \quad (3.4)$$

It can easily be checked that identities (3.3) are fulfilled if the condition $\{\tilde{H}, \tilde{K}\} = 0$ holds, where

$$\tilde{K} = -(s_1 - s_2)^2 p_1 p_2.$$

It is easy to verify that if $\{p_i, s_j\} = \delta_{ij}$, then the Poisson brackets between functions

$$a_0 = p_1 + p_2, \quad a_1 = s_1 p_1 + s_2 p_2, \quad a_2 = s_1^2 p_1 + s_2^2 p_2 \quad (3.5)$$

are given by

$$\{a_0, a_1\} = a_0, \quad \{a_0, a_2\} = 2a_1, \quad \{a_1, a_2\} = a_2. \quad (3.6)$$

The expression $Q = a_1^2 - a_0 a_2$ is a Casimir function for the linear $sl(2)$ -brackets (3.6). It turns out that \tilde{K} rewritten in variables (3.5) coincides with Q . The Hamiltonian \tilde{H} can also be expressed in terms of variables (3.5) only:

$$\tilde{H} = a_0^3 L \left(\frac{a_1}{a_0} \right) - \delta^2 (\alpha a_2 + e_2 a_1 + e_1 a_0) + \frac{\delta^4}{4} \frac{a_1^2}{a_0}.$$

It is seen from this formula that the variable

$$\xi = \frac{a_1}{a_0} = \frac{p_1 s_1 + p_2 s_2}{p_1 + p_2}$$

should play a key role in a description of properties of polynomials (2.10), (3.4). Suppose $\tilde{H}(p_1, p_2) = 0$; then a solution z for equation (2.10) can be expressed in terms of ξ by the formula

$$z = \sqrt{\frac{s_2 - \xi}{s_1 - \xi}} \sqrt{\frac{S(s_1)}{S(s_2)}}. \quad (3.7)$$

Equation (2.10) is cubic with respect to ξ . It can be rewritten as (1.1), where

$$Y^2 \stackrel{\text{def}}{=} \frac{1}{\delta^2} (\xi - s_1)(\xi - s_2). \quad (3.8)$$

Our task is to find the solution of (3.2) explicitly. According to formulas (2.11), it suffices for that to solve the system of equation

$$\begin{aligned} \frac{\partial \sigma}{\partial s_1} &= \frac{J}{8S(s_1)} z = \frac{\delta}{4(s_1 - s_2)} \sqrt{\frac{s_2 - \xi}{s_1 - \xi}}, \\ \frac{\partial \sigma}{\partial s_2} &= -\frac{J}{8S(s_2)} \frac{1}{z} = -\frac{\delta}{4(s_1 - s_2)} \sqrt{\frac{s_1 - \xi}{s_2 - \xi}}. \end{aligned} \quad (3.9)$$

Here $z(s_1, s_2)$ is arbitrary root of equation (2.10) and ξ is the corresponding (see (3.7)) value of $\xi(s_1, s_2)$. The action function \tilde{S} required can be obtained as the sum of the three solutions for (3.9), corresponding to the three branches of the function $\xi(s_1, s_2)$.

First, we find a function σ_0 such that (3.9) is fulfilled under condition that ξ is a parameter, which does not depend on s_1, s_2 . It is not hard to see that

$$\sigma_0(\xi, s_1, s_2) = \frac{1}{4} \delta \operatorname{arctanh} \frac{\xi - \frac{1}{2}(s_1 + s_2)}{\sqrt{(\xi - s_1)(\xi - s_2)}}$$

Taking into account the formula

$$\frac{\partial}{\partial \xi} \sigma_0 = \frac{1}{4} \frac{1}{\sqrt{(\xi - s_1)(\xi - s_2)}}$$

and replacing $\sqrt{(\xi - s_1)(\xi - s_2)}$ by $Y\delta$, we obtain expression (3.1) for the action function.

Differentiating the action with respect to the parameters and taking into account that

$$\frac{\partial \tilde{S}}{\partial e_1} = t + c_1, \quad \frac{\partial \tilde{S}}{\partial e_2} = c_2,$$

we finally get

$$dt = \sum_{n=1}^3 \omega_1(\xi_n), \quad 0 = \sum_{n=1}^3 \omega_2(\xi_n). \quad (3.10)$$

Here ω_1, ω_2 belong to a basis

$$\omega_1(\xi) = \frac{d\xi}{Z}, \quad \omega_2(\xi) = \frac{\xi d\xi}{Z}, \quad \omega_3(\xi) = \frac{(\xi^2 - \delta^2 Y^2) d\xi}{Z}, \quad \omega_4(\xi) = \frac{Y d\xi}{Z} \quad (3.11)$$

of holomorphic differentials on the curve $\Phi(\xi, Y) = 0$. In (3.11) we use the notation $Z \stackrel{\text{def}}{=} \frac{\partial \Phi}{\partial Y}$.

The differential ω_4 plays a special role. In variables (ξ, η) , where

$$\eta = \xi^2 - \delta^2 Y^2, \quad (3.12)$$

the curve (1.1) becomes the following cubic

$$s_6 \eta^3 + s_5 \eta^2 \xi + \frac{1}{5} s_4 \eta (\eta + 4\xi^2) + \frac{1}{5} s_3 \xi (\eta + 2\xi^2) + \frac{1}{5} s_2 (\eta + 4\xi^2) + s_1 \xi + s_0 + \frac{4}{\delta^2} (\eta - \xi^2) (\alpha \eta + e_2 \xi + e_1) = 0. \quad (3.13)$$

The holomorphic differential of the elliptic curve (3.13) coincides with ω_4 up to transformation (3.12).

It follows from (3.8) that the functions $Y(\xi_i)$ are linked by the relation

$$\frac{1}{\delta^2} = \frac{Y^2(\xi_1)}{(\xi_1 - \xi_2)(\xi_1 - \xi_3)} + \frac{Y^2(\xi_2)}{(\xi_2 - \xi_3)(\xi_2 - \xi_1)} + \frac{Y^2(\xi_3)}{(\xi_3 - \xi_1)(\xi_3 - \xi_2)}, \quad (3.14)$$

which allows us to determine the two functions $s_1(t), s_2(t)$ starting from $\xi_1(t), \xi_2(t), \xi_3(t)$. Thus we have three conditions (3.10), (3.14) to determine three functions $\xi_i(t)$.

Condition (3.14) can be rewritten in the variables (ξ, η) as

$$\eta_1(\xi_2 - \xi_3) + \eta_2(\xi_3 - \xi_1) + \eta_3(\xi_1 - \xi_2) = 0.$$

This means that the corresponding points (ξ_i, η_i) belongs to the intersection of elliptic curve (3.13) and the straight line $\eta = \xi(s_1 + s_2) - s_1 s_2$.

4 Schottky-Manakov spinning top.

It is well-known (see, for example [19, 11]) that the Hamiltonian

$$H = (\vec{S}_1, A\vec{S}_1) + 2(\vec{S}_1, B\vec{S}_2) + (\vec{S}_2, A\vec{S}_2),$$

where $A = \text{diag}(a_1, a_2, a_3)$, $B = \text{diag}(b_1, b_2, b_3)$, commutes with a quadratic polynomial K with respect to the spin Poisson brackets

$$\{S_i^\alpha, S_j^\beta\} = \kappa \varepsilon_{\alpha\beta\gamma} S_i^\gamma \delta_{ij} \quad (4.1)$$

iff

$$b_1^2(a_2 - a_3) + b_2^2(a_3 - a_1) + b_3^2(a_1 - a_2) + (a_1 - a_2)(a_2 - a_3)(a_3 - a_1) = 0. \quad (4.2)$$

Here $\varepsilon_{\alpha\beta\gamma}$ is the totally skew-symmetric tensor, κ is a parameter.

Since H and K may be replaced by arbitrary linear combinations of H , K and the Casimir functions

$$J_1 = (\vec{S}_1, \vec{S}_1), \quad J_2 = (\vec{S}_2, \vec{S}_2)$$

for the brackets (4.1), the integral K can be reduced [13] to the form

$$K = 2(\vec{S}_1, \hat{C}\vec{S}_2), \quad C = \text{diag}(\alpha_1, \alpha_2, \alpha_3).$$

Without loss of generality the matrices A and B defined the Hamiltonian H can be choosen as follows

$$A = \text{diag}(-\alpha_1^2, -\alpha_2^2, -\alpha_3^2), \quad B = \text{diag}(\alpha_2\alpha_3 + \lambda\alpha_1, \alpha_3\alpha_1 + \lambda\alpha_2, \alpha_1\alpha_2 + \lambda\alpha_3),$$

The arbitrary parameter λ corresponds to the shift of H by λK .

Reduction to the standard brackets. Let us fix the values of the Casimir functions: $(\vec{S}_k, \vec{S}_k) = j_k^2$. Then the formulas

$$\vec{S}_k = p_k \vec{K}(q_k) + \frac{j_k}{2} \vec{K}'(q_k), \quad \text{where} \quad \vec{K}(q) = ((q^2 - 1), i(q^2 + 1), 2q), \quad (4.3)$$

define a transformation of the Poisson manifold with coordinates \vec{S}_1, \vec{S}_2 and brackets (4.1), where $\kappa = -2i$, to the manifold with coordinates p_1, p_2, q_1, q_2 and canonical Poisson brackets $\{p_\alpha, q_\beta\} = \delta_{\alpha\beta}$. As a result of this transformation we get

$$\begin{aligned} H = & p_1^2 r(q_1) + \frac{j_1}{2} p_1 r'(q_1) + \frac{j_1^2}{12} r''(q_1) + p_2^2 r(q_2) + \frac{j_2}{2} p_2 r'(q_2) + \frac{j_2^2}{12} r''(q_2) + \\ & 2 \left(p_1 + \frac{j_1}{2} \frac{\partial}{\partial q_1} \right) \left(p_2 + \frac{j_2}{2} \frac{\partial}{\partial q_2} \right) Z(q_1, q_2), \end{aligned} \quad (4.4)$$

$$K = 2 \left(p_1 + \frac{j_1}{2} \frac{\partial}{\partial q_1} \right) \left(p_2 + \frac{j_2}{2} \frac{\partial}{\partial q_2} \right) W(q_1, q_2), \quad (4.5)$$

where

$$\begin{aligned} r(x) &\stackrel{\text{def}}{=} (\vec{K}(x), A\vec{K}(x)) = -\alpha_1^2(x^2 - 1)^2 + \alpha_2(x^2 + 1)^2 - 4\alpha_3x^2, \\ Z(x, y) &\stackrel{\text{def}}{=} (\vec{K}(x), B\vec{K}(y)) = \\ &(\alpha_2\alpha_3 + \lambda\alpha_1)(x^2 - 1)(y^2 - 1) - (\alpha_3\alpha_1 + \lambda\alpha_2)(x^2 + 1)(y^2 + 1) + 4(\alpha_1\alpha_2 + \lambda\alpha_3)xy, \end{aligned} \quad (4.6)$$

$$W(x, y) \stackrel{\text{def}}{=} (\vec{K}(x), C\vec{K}(y)) = \alpha_1(x^2 - 1)(y^2 - 1) - \alpha_2(x^2 + 1)(y^2 + 1) + 4\alpha_3xy. \quad (4.7)$$

It is possible to check that

$$Z^2(x, y) - r(x)r(y) = W(x, y)\bar{W}(x, y), \quad (4.8)$$

where \bar{W} is a polynomial quadratic in each of variables.

Remark. Note that the more general Hamiltonian

$$H = \sum_{ij, \mu, \nu} S_i^\mu S_j^\nu c_{\mu\nu}^{ij}, \quad i, j = 1, \dots, N, \quad \mu, \nu = x, y, z,$$

describing an interaction of N spins, can be reduced to

$$H = \sum_{ij} g^{ij} p_i p_j + \sum_i a^i p_i + v,$$

where

$$\begin{aligned} g^{ij} &= \sum_{\mu, \nu} K^\mu(x^i) K^\nu(x^j) c_{\mu\nu}^{ij}, \\ a^i &= \sum_k j_k \frac{\partial g^{ik}}{\partial x^k} - \frac{1}{2} j_i \frac{\partial g^{ii}}{\partial x^i}, \\ v &= \frac{1}{4} \sum_{ik} j_i j_k \frac{\partial^2 g^{ik}}{\partial x^i \partial x^k} - \frac{1}{6} \sum_k j_k^2 \frac{\partial^2 g^{kk}}{\partial x^k \partial x^k} \end{aligned}$$

by a similar transformation

$$\vec{S}_k = p_k \vec{K}(x^k) + \frac{j_k}{2} \vec{K}'(x^k), \quad k = 1, \dots, N.$$

In terms of the coordinates and the velocities the Lagrangian and the energy read as

$$L = \frac{1}{4} \sum_{ij} g_{ij} (\dot{x}^i - a^i)(\dot{x}^j - a^j) - v, \quad E = \frac{1}{4} \sum_{ij} g_{ij} (\dot{x}^i \dot{x}^j - a^i a^j) + v,$$

where $\sum_j g_{ij} g^{jk} = \delta_i^k$.

An additional integral of the form

$$K = \sum_{ij} G^{ij} p_i p_j + \sum_i A^i p_i + V$$

exists iff

$$\sum_k \frac{\partial G^{ij}}{\partial x^k} g^{kl} - \frac{\partial g^{ij}}{\partial x^k} G^{kl} = 0, \quad \sum_k a^k \frac{\partial G^{ij}}{\partial x^k} - 2 \frac{\partial a^i}{\partial x^k} G^{kj} - A^k \frac{\partial g^{ij}}{\partial x^k} + 2 \frac{\partial A^i}{\partial x^k} g^{kj} = 0,$$

$$\sum_k a^k \frac{\partial A^i}{\partial x^k} - A^k \frac{\partial a^i}{\partial x^k} + 2g^{ik} \frac{\partial V}{\partial x^k} - 2G^{ik} \frac{\partial v}{\partial x^k} = 0, \quad \sum_k a^k \frac{\partial V}{\partial x^k} - A^k \frac{\partial v}{\partial x^k} = 0.$$

Diagonalization of the quadratic part. Under transformation (4.3) the functions H and K take the form

$$H = ap_1^2 + 2bp_1p_2 + cp_2^2 + dp_1 + ep_2 + f, \quad (4.9)$$

$$K = Ap_1^2 + 2Bp_1p_2 + Cp_2^2 + Dp_1 + Ep_2 + F, \quad (4.10)$$

where the coefficients are some (in our case rational) functions of the variables q_1, q_2 .

General formulas related to a pair of commuting Hamiltonians quadratic in momenta and some examples can be found in [16, 14, 15, 17].

The class of Hamiltonians (4.9) is invariant with respect to *canoniacal* transformations of the form

$$p_1 = k_1 \hat{p}_1 + k_2 \hat{p}_2 + k_3, \quad p_2 = \bar{k}_1 \hat{p}_1 + \bar{k}_2 \hat{p}_2 + \bar{k}_3, \quad q_1 = \phi, \quad q_2 = \bar{\phi},$$

where $k_i, \bar{k}_i, \phi, \bar{\phi}$ are some functions of \hat{q}_1, \hat{q}_2 . It is easily seen that the functions $k_1, k_2, \bar{k}_1, \bar{k}_2$ can be uniquely expressed through $\phi, \bar{\phi}$ from the condition $\{p_\alpha, q_\beta\} = \delta_{\alpha\beta}$:

$$k_1 = \frac{\partial \bar{\phi}}{\partial \hat{q}_2} W^{-1}, \quad k_2 = -\frac{\partial \bar{\phi}}{\partial \hat{q}_1} W^{-1}, \quad \bar{k}_1 = -\frac{\partial \phi}{\partial \hat{q}_2} W^{-1}, \quad \bar{k}_2 = \frac{\partial \phi}{\partial \hat{q}_1},$$

where

$$W = \frac{\partial \bar{\phi}}{\partial \hat{q}_2} \frac{\partial \phi}{\partial \hat{q}_1} - \frac{\partial \bar{\phi}}{\partial \hat{q}_1} \frac{\partial \phi}{\partial \hat{q}_2}.$$

The fact that the coefficient of $\hat{p}_1 \hat{p}_2$ in the transformed Hamiltonian (4.9) vanishes is equivalent to

$$a(\phi, \bar{\phi}) \frac{\partial \bar{\phi}}{\partial \hat{q}_1} \frac{\partial \bar{\phi}}{\partial \hat{q}_2} + c(\phi, \bar{\phi}) \frac{\partial \phi}{\partial \hat{q}_1} \frac{\partial \phi}{\partial \hat{q}_2} = b(\phi, \bar{\phi}) \left(\frac{\partial \bar{\phi}}{\partial \hat{q}_2} \frac{\partial \phi}{\partial \hat{q}_1} + \frac{\partial \bar{\phi}}{\partial \hat{q}_1} \frac{\partial \phi}{\partial \hat{q}_2} \right). \quad (4.11)$$

Analogously, the condition

$$A(\phi, \bar{\phi}) \frac{\partial \bar{\phi}}{\partial \hat{q}_1} \frac{\partial \bar{\phi}}{\partial \hat{q}_2} + C(\phi, \bar{\phi}) \frac{\partial \phi}{\partial \hat{q}_1} \frac{\partial \phi}{\partial \hat{q}_2} = B(\phi, \bar{\phi}) \left(\frac{\partial \bar{\phi}}{\partial \hat{q}_2} \frac{\partial \phi}{\partial \hat{q}_1} + \frac{\partial \bar{\phi}}{\partial \hat{q}_1} \frac{\partial \phi}{\partial \hat{q}_2} \right) \quad (4.12)$$

guarantees that the coefficient of $\hat{p}_1\hat{p}_2$ in the transformed integral (4.10) is equal to zero. The existence of a solution $\phi, \bar{\phi}$ for system (4.11), (4.12) is obvious. Our goal is to find the functions $\phi, \bar{\phi}$ for the pair (4.4), (4.5) explicitly.

Let us introduce the Kowalewski variables $s_1(q_1, q_2)$ and $s_2(q_1, q_2)$ as the roots of the equation

$$W(q_1, q_2) s^2 + 2Z(q_1, q_2) s + \bar{W}(q_1, q_2) = 0,$$

where W and Z are given by (4.7), (4.6) and the polynomial \bar{W} is defined by (4.8).

Proposition 3. *In the variables s_1, s_2 the functions H and K have the form (1.3)-(1.5), where polynomials S and R are defined by the formulas*

$$S(x) = -(x + \lambda - \alpha_1 - \alpha_2 - \alpha_3)(x + \lambda + \alpha_1 + \alpha_2 - \alpha_3)(x + \lambda + \alpha_1 - \alpha_2 + \alpha_3)(x + \lambda - \alpha_1 + \alpha_2 + \alpha_3)$$

and

$$\alpha = \frac{1}{4}\left(\frac{\delta^2}{5} - \nu^2\right), \quad \delta = j_1 - j_2, \quad \nu = j_1 + j_2.$$

The values h and k of the integrals H and K are related to the constants e_1, e_2 from (1.2) by

$$h = e_1 + \frac{1}{12}\alpha S''(0), \quad k = e_2 + \frac{1}{12}\alpha S'''(0).$$

Thus the general scheme from Section 3 can be applied to the Schottky-Manakov spinning top.

5 The Kowalewski gyrostat.

In this Section we use some formulas from [21] that describe a dynamical system for the Kowalewski gyrostat in the Kowalewski variables.

The Hamiltonian structure for the gyrostat is defined by the $e(3)$ -Poisson brackets

$$\{M_i, M_j\} = \varepsilon_{ijk} M_k, \quad \{M_i, \gamma_j\} = \varepsilon_{ijk} \gamma_k, \quad \{\gamma_i, \gamma_j\} = 0,$$

where ε_{ijk} is the totally skew-symmetric tensor. The brackets possess the Casimir functions

$$A = \sum_{k=1}^3 \gamma_k^2, \quad B = \sum_{k=1}^3 \gamma_k M_k. \quad (5.1)$$

The Hamiltonian for the gyrostat is as follows:

$$H = \frac{1}{2}(M_1^2 + M_2^2 + 2M_3^2 - 2\lambda M_3) + c\gamma_1, \quad (5.2)$$

where c and λ are constants. An additional integral of motion is given by

$$K = \xi_1 \xi_2 + 4\lambda \left((M_3 - \lambda) z_1 z_2 - (z_1 + z_2) c \gamma_3 \right), \quad (5.3)$$

where

$$\xi_1 = z_1^2 - 2c(\gamma_1 + i\gamma_2), \quad \xi_2 = z_2^2 - 2c(\gamma_1 - i\gamma_2)$$

and

$$z_1 = M_1 + iM_2, \quad z_2 = M_1 - iM_2.$$

We put

$$R(z_1, z_2) = z_1^2 z_2^2 - 2h(z_1^2 + z_2^2) - 4cb(z_1 + z_2) - 4c^2 a + k.$$

Here a , b , h and k are values of integrals (5.1), (5.2), and (5.3). The Kowalewski variables are defined as follows

$$s_{1,2} = \frac{R(z_1, z_2) \pm \sqrt{R(z_1, z_1)R(z_2, z_2)}}{2(z_1 - z_2)^2}.$$

It turns out [21] that

$$h = \frac{s_1 - s_2}{2} \left(\frac{\dot{s}_1^2}{\varphi_1} - \frac{\dot{s}_2^2}{\varphi_2} \right) - \frac{s_1 + s_2}{2}, \quad (5.4)$$

$$\frac{k}{4} = (2h + s_1 + s_2)\lambda^2 - \lambda\sqrt{-\varphi_1\varphi_2} \left(\frac{\dot{s}_1}{\varphi_1} + \frac{\dot{s}_2}{\varphi_2} \right) + (s_1 - s_2) \left(\frac{s_2\dot{s}_1^2}{\varphi_1} - \frac{s_1\dot{s}_2^2}{\varphi_2} \right) - s_1 s_2 + h^2. \quad (5.5)$$

Here $\varphi_i = S(s_i)$,

$$S(s) = 4s^3 - 8hs^2 + 4h^2s - ks + 4c^2as + 4c^2b.$$

Substituting the expressions (2.8)

$$\dot{s}_1 = -\frac{1}{4} \frac{J}{s_1 - s_2} (u + 1), \quad \dot{s}_2 = -\frac{1}{4} \frac{J}{s_1 - s_2} (v + 1)$$

in (5.4), (5.5) for the velocities, we get just (2.5), where

$$k(x) = 4(x + h)^2 + 4\delta^2(x - 2h) - k, \quad \lambda = i\delta.$$

Thus in this case the scheme for the separation of variables described in Section 3 is applicable. The curve $\Phi(Y, \xi) = 0$ is of genus 3. The differentials $\omega_1, \omega_2, \omega_4$ form a basis of holomorphic differentials.

Notice that the substitution $\delta = -i\lambda$ in (3.1) results the change $\arctanh \rightarrow \arctan$ and the action remains to be real.

6 The Clebsch spinning top.

The Clebsch spinning top is defined by the Hamiltonian

$$H = \frac{1}{2}(J_1^2 + J_2^2 + J_3^2) + \frac{1}{2}(\lambda_1 x_1^2 + \lambda_2 x_2^2 + \lambda_3 x_3^2)$$

which commutes with respect to the $e(3)$ -Poisson brackets

$$\{J_i, J_j\} = i \varepsilon_{ijk} J_k, \quad \{x_i, x_j\} = 0, \quad \{J_i, x_j\} = i \varepsilon_{ijk} x_k$$

with the first integral

$$K = (\lambda_1 J_1^2 + \lambda_2 J_2^2 + \lambda_3 J_3^2) - \lambda_1 \lambda_2 \lambda_3 \left(\frac{x_1^2}{\lambda_1} + \frac{x_2^2}{\lambda_2} + \frac{x_3^2}{\lambda_3} \right).$$

Let us fix the values of the Casimir functions as follows

$$x_1^2 + x_2^2 + x_3^2 = a^2, \quad J_1 x_1 + J_2 x_2 + J_3 x_3 = l.$$

Using the parameterization

$$J_1 = \frac{1}{2}p_1(1-q_1^2) + \frac{1}{2}p_2(1-q_2^2) + \frac{l}{a}q_1, \quad J_2 = \frac{i}{2}p_1(1+q_1^2) + \frac{i}{2}p_2(1+q_2^2) - i\frac{l}{a}q_1, \quad J_3 = p_1 q_1 + p_2 q_2 - \frac{l}{a},$$

$$x_1 = a \frac{1 - q_1 q_2}{q_1 - q_2}, \quad x_2 = i a \frac{1 + q_1 q_2}{q_1 - q_2}, \quad x_3 = a \frac{q_1 + q_2}{q_1 - q_2},$$

we express H and K in terms of canonically conjugated variables p_1, q_1, p_2, q_2 :

$$H = -\frac{1}{2}(x-y)^2 p_1 p_2 + \frac{l}{a} p_2 (x-y) + \frac{a^2}{2} \frac{W(x,y)}{(x-y)^2} + \frac{1}{2}(\lambda a^2 + \frac{l^2}{a^2})$$

$$K = \frac{1}{4}(R(x)p_1^2 + R(y)p_2^2 + 2p_1 p_2 W(x,y)) - \frac{l}{4a} p_1 R'(x) - \frac{l}{2a} p_2 W_x(x,y) - a^2 \frac{\bar{W}(x,y)}{(x-y)^2} + \frac{l^2 R''(x)}{12a^2} + \frac{\lambda l^2}{3a^2}$$

where

$$R(x) = (\lambda_1 - \lambda_2)(x^2 - 1)^2 + 4(\lambda_3 - \lambda_2)x^2,$$

$$W(x,y) = \lambda_1(x^2 - 1)(y^2 - 1) - \lambda_2(x^2 + 1)(y^2 + 1) + 4\lambda_3 xy,$$

$$\bar{W}(x,y) = \lambda_2 \lambda_3 (x^2 - 1)(y^2 - 1) - \lambda_3 \lambda_1 (x^2 + 1)(y^2 + 1) + 4\lambda_1 \lambda_2 xy + \kappa(x-y)^2,$$

$$\lambda = \lambda_1 + \lambda_2 + \lambda_3, \quad \kappa = \lambda_1 \lambda_2 + \lambda_2 \lambda_3 + \lambda_3 \lambda_1.$$

We define Kowalewski variables as the roots $s_1(q_1, q_2)$ and $s_2(q_1, q_2)$ of the equation

$$(q_1 - q_2)^2 s^2 + W(q_1, q_2) s + \bar{W}(q_1, q_2) = 0.$$

Associating the t -dynamics with K , we get equations (2.3), (2.4) for s_1, s_2 , where

$$S(\xi) = 4(\xi - \lambda_1)(\xi - \lambda_2)(\xi - \lambda_3).$$

Our general procedure for separation of variables from Section 3 leads to curve (1.1), where

$$s_6 = 0, \quad l(\xi) = \frac{l^2}{64}, \quad 4k(\xi) = a^2\xi^2 + (2e_1 - a^2\lambda)\xi - e_2.$$

Note that for the Clebsch case we have $\delta = i\frac{l}{4a}$. One has to replace $\operatorname{arctanh}$ by arctan in the formula (3.1) and after that the action remains to be real.

7 Case $S = -1$

In the case $S(\xi) = -1$ common solutions to the pair (1.3), (1.5) satisfy the Gibbons-Tsarev equation (2.7). In this case the curve (1.1) has the form

$$-4\alpha\delta^2Y^4 + (4e_1 + 4\xi e_2 + 4\xi^2\alpha)Y^2 + 1 = 0.$$

This curve of genus 1 can be parameterized by the Weierstrass function as follows:

$$Y = -\frac{1}{16\alpha^2} \frac{y}{x - \beta_3}, \quad \xi = -\frac{e_2}{2\alpha} - \frac{y}{16\alpha^2} \left(\frac{1}{x - \beta_1} + \frac{1}{x - \beta_2} - \frac{1}{x - \beta_3} \right),$$

where $x = \wp(z, g_2, g_3)$, $y = \wp'(z, g_2, g_3)$,

$$y^2 = 4(x - \beta_1)(x - \beta_2)(x - \beta_3), \quad \beta_3 = \frac{8\alpha^2}{3\delta^2}(4\alpha e_1 - e_2^2), \quad \beta_{1,2} = -\frac{1}{2}\beta_3 \pm \frac{16\alpha^3}{\delta}\sqrt{-\alpha}.$$

The meromorphic integrals in

$$dt = \sum_{n=1}^3 \omega_1(\xi_n), \quad d\tau = \sum_{n=1}^3 \omega_2(\xi_n)$$

can be taken. The result is given by

$$z_1 + z_2 + z_3 = \text{const}, \quad t = \frac{1}{2\sqrt{-\alpha}} \sum_{i=1}^3 \log\left(\frac{x_i - \beta_2}{\beta_1 - x_i}\right), \quad \tau = -\frac{e_2}{\alpha}t - 2\alpha \sum_{i=1}^3 \frac{y_i}{(x_i - \beta_1)(x_i - \beta_2)}$$

where $x_i = \wp(z_i, g_2, g_3)$, $y_i = \wp'(z_i, g_2, g_3)$.

References

- [1] F Schottky, Über das analytische Problem der Rotation eines starren Körpers in Raume von vier Dimensionen. *Sitzungsberichte drer Königlich preussischen Academie der Wissenschaften zu Berlin*, **XIII**, 227–232, 1891.
- [2] S V Manakov, A remark on integration of the Euler equations for n -dimensional rigid body dynamics, *Funct. Anal. Appl.*, **10**(4)), 93–94, 1976.
- [3] Yu N Fedorov, Integrable systems, Lax representations, and confocal quadrics. Dynamical systems in classical mechanics, 173–199, Amer. Math. Soc. Transl. Ser. 2, **168**, Amer. Math. Soc., Providence, RI, 1995.
- [4] A Clebsch, Über die Bewegung eines Körpers in einer Flüssigkeit, *Math. Annalen*, **3**, 238–262, 1870.
- [5] A I Perelomov, Some remarks on the integrability of the equations of motion of a rigid body in an ideal fluid, *Funct. Anal. Appl.*, **15**, 83–85, 1981.
- [6] F Kötter, Über die Bewegung eines festen Körpers in einer Flüssigkeit. I.II. *J. Reine und Angew. Math.*, **109**, 51–81, 89–111, 1892.
- [7] A G Reyman and M A Semenov-tian-Shanski, Integrable systems, *Moscow-Izevsk*, 352 p., 2003.
- [8] A V Borisov and S I Mamaev, Modern methods of the theory of integrable systems, *Moscow-Izevsk*, 296 p., 2003.
- [9] J Gibbons and S P Tsarev, Reductions of the Benney equations, *Phys. Lett. A*, **211**, 19–24, 1996.
- [10] L Haine and E Horozov, A Lax pair for the Kowalewski top, *Physica D.*, **29**, 173–180, 1987.
- [11] A S Mischenko, Integration of geodesic flows on the symmetric spaces, *Math. Notes*, **31**(2)), 257–262, 1982.
- [12] A S Mischenko and A T Fomenko, Euler equations on finite-dimensional Lie groups, *Izvestiya Akad. Nauk SSSR, ser. math.*, **42**(2)), 396–415, 1978.
- [13] E G Kalnins, W (JR) Miller, and P Winternitz, The group $O(4)$, separation of variables and the hydrogen atom, *SIAM J. Appl. Math.*, 1976, v.30, 4, p.630–664.

- [14] E V Ferapontov and A P Fordy, Nonhomogeneous systems of hydrodynamic type related to quadratic Hamiltonians with electromagnetic term, *Physica D* 108 (1997) 350-364.
- [15] E V Ferapontov and A P Fordy, Commuting quadratic Hamiltonians with velocity-dependent potentials, *Rep. Math. Phys.* 44, N1/2 (1999) 71-80.
- [16] B Dorizzi, B Grammaticos, A Ramani and P Winternitz, Integrable Hamiltonian systems with velocity dependent potentials, *J Math. Phys.* 26 (1985) 3070-3079.
- [17] E McSween and P Winternitz, Integrable and superintegrable Hamiltonian systems in magnetic fields, *J Math Phys* 41 (2000) 2957-2967.
- [18] M Adler and P van Moerbeke. The Kowalewski and Henon-Heiles motions as Manakov geodesic flows on $so(4)$ - a two dimensional family of Lax pairs, *Comm. in Math. Phys.*, **113**, 659–700, 1988.
- [19] A P Veselov, On the integrability conditions for the Euler equation on $so(4)$, *Dokl. Akad. Nauk SSSR*, **270**(6), 1298–1300, 1983.
- [20] A I Bobenko, The Euler equations on $e(3)$ and $so(4)$. Isomorphism of integrable cases. *Funct. Anal. Appl.* **20**(1), 64–66, 1986.
- [21] I V Komarov and A V Tsiganov, On integration of the Kowalewski gyrostat problem, *Reg. Chaot. Dyn.*, **9**(2), 169-187, 2004.